

Expanding the Oro-Dental and Mutational Spectra of Kabuki Syndrome and

Expression of *KMT2D* and *KDM6A* in Human Tooth Germs

Thanrira Porntaveetus¹, Mushriq F Abid², Thanakorn Theerapanon³, Chalurmpon Srichomthong^{4,5}, Atsushi Ohazama⁶, Katsushige Kawasaki⁶, Maiko Kawasaki⁶, Kanya Suphapeetiporn^{4,5}, Paul T. Sharpe², Vorasuk Shotelersuk^{4,5}

Table S1. Primer sequences

For sanger sequencing			
Gene	Exon	Forward	Reverse
<i>KMT2D</i>	7	CGG AGA AGG CCT GTT AAC AT	AGT TGG GAT TCA GTT CTG CT
<i>KMT2D</i>	11	CAC ACC ACC CAC TCT CAT CA	AAT CAC ATC CCG CAG CTA GA
<i>KMT2D</i>	22	TGA CAC TTC CGT TCT GTC CA	ACT GCA AAA AGG GCC TTA CG
<i>KMT2D</i>	29	ACC AAC AGC AAG GGT CCA T	GTG TTT TGT TCC TTG CCC GT
<i>KMT2D</i>	39	GCT GGG AGA AGG ATG AGG AG	CCT ACC CAG AAG CAG ATC CC
<i>KMT2D</i>	51	AA CAA CGT GTA CCT GGC TC	CAC AGC TTT CCT CCT GCT CT
<i>KDM6A</i>	3	TAA ATA CAT AGG GAA GCT GTT C	AAT CCA ACA CAT GTA TAA ATC AAA C

KMT2D, Lysine-specific methyltransferase 2D; *KDM6A*, Lysine-specific demethylase 6A