

2024; 20(15): 6038-6055. doi: 10.7150/ijbs.103977

Review

# Multi-Omics Approaches to Discover Biomarkers of Thyroid Eye Disease: A Systematic Review

Haiyang Zhang<sup>1,2,3,†</sup>, Yuyu Zhou<sup>1,2,3,†</sup>, Baiguang Yu<sup>1,2,3,†</sup>, Yuyang Deng<sup>1,2,3</sup>, Yang Wang<sup>1,2,3</sup>, Sijie Fang<sup>1,2,3</sup>, Xuefei Song<sup>1,2,3,\infty</sup>, Xianqun Fan<sup>1,2,3,\infty</sup>, Huifang Zhou<sup>1,2,3,\infty</sup>

- 1. Department of Ophthalmology, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, No.639, Zhizaoju Road, Huangpu District, Shanghai, China.
- 2. Shanghai Key Laboratory of Orbital Diseases and Ocular Oncology, Shanghai, China.
- 3. Center for Basic Medical Research and Innovation in Visual System Diseases, Ministry of Education, Shanghai, China.

†Haiyang Zhang, Yuyu Zhou, and Baiguang Yu contributed equally to this manuscript.

🖂 Corresponding authors: Huifang Zhou, fangzzfang@sjtu.edu.cn; Xianqun Fan, fanxq@sjtu.edu.cn; Xuefei Song, songxuefei@shsmu.edu.cn.

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/). See https://ivyspring.com/terms for full terms and conditions.

Received: 2024.09.21; Accepted: 2024.10.28; Published: 2024.11.11

#### Abstract

Thyroid eye disease (TED) is an organ-specific autoimmune disorder that significantly impacts patients' visual function, appearance, and well-being. Despite existing clinical evaluation methods, there remains a need for objective biomarkers to facilitate clinical management and pathogenesis investigation. Rapid advances in multi-omics technologies have enabled the discovery and development of more informative biomarkers for clinical use. This systematic review synthesizes the current landscape of multi-omics approaches in TED research, highlighting the potential of genomics, transcriptomics, proteomics, metabolomics, and microbiomics to uncover novel biomarkers. Our review encompasses 69 studies involving 1,363 TED patients and 1,504 controls, revealing a wealth of biomarker candidates across various biological matrices. The identified biomarkers reflect alterations in gene expression, protein profiles, metabolic pathways, and microbial compositions, underscoring the systemic nature of TED. Notably, the integration of multi-omics data has been pivotal in enhancing our understanding of TED's molecular mechanisms and identifying diagnostic and prognostic markers with clinical potential.

Keywords: Thyroid eye disease, Biomarker, Multi-omics, Bioinformation

## Introduction

Thyroid eve disease (TED), also known as thyroid-associated ophthalmopathy (TAO) and Graves' orbitopathy (GO), is an organ-specific autoimmune disorder characterized bv the enlargement of the extraocular muscles and an increase in fatty or connective tissue volume. It is the most frequent extrathyroidal manifestation of Graves' disease (GD) and occasionally occurs in patients with autoimmune thyroiditis and other thyroid diseases, resulting from a combination of genetic and environmental factors [1]. The estimated incidence rate is 4.2 per 100,000 person-years. Female patients outnumber males, and it tends to occur in individuals aged 40-50 years and 60-70 years [2]. The clinical manifestations of TED include upper evelid retraction, oedema, erythema of the periorbital tissues

and conjunctivae, exophthalmos, dry eye, diplopia, and strabismus, significantly affecting the quality of life of patients [3, 4].

The pathogenesis of TED is not vet fully but autoantibodies understood, to the thyroid-stimulating hormone receptor (TSH-R) and insulin-like growth factor-1 receptor (IGF-1R) are thought to play a key role [4, 5]. Current disease activity scoring relies on the European Group on Graves' Orbitopathy (EUGOGO) system for clinical activity scores (CAS), which considers the presence of pain, redness, swelling, and functional impairment of specific ocular and orbital structures [6, 7]. However, the pathological processes of TED extend beyond orbital inflammation and tissue remodeling to include immune and metabolic changes in peripheral blood

[8–11]. These changes serve as important biomarkers for a deeper understanding of the disease's complexity. For instance, intraorbital biomarkers may describe the activation of orbital fibroblasts, changes in the fibrosis of extraocular muscles, and the proliferation of adipose tissues [12]. In peripheral blood, biomarkers related to inflammation, immune cell activation, and autoimmune responses may be observed, such elevated levels as of thyroid-stimulating hormone receptor antibodies (TRAb), the presence of specific antibodies, or alterations in the proportions of immune cell subsets [13]. These biomarkers are valuable not only for early diagnosis and assessment of disease activity but also for predicting disease progression and treatment response. Therefore, identifying and validating these biomarkers is crucial.

With the rapid development of high-throughput sequencing technology and the popularization of public databases, multi-omics studies using bioinformatics analysis have been conducted to identify new diagnostic or therapeutic biomarkers for various diseases, including cancers and autoimmune disorders [14, 15]. According to the central dogma of molecular biology, omics fields such as genomics, transcriptomics, proteomics, epigenomics, metabolomics, and lipidomics have a promising role biomarkers in investigating more and comprehensively understanding the molecular mechanisms of TED [16]. For example, several microarray gene profile studies on TED have been performed to identify differentially expressed genes (DEGs) and key pathways, revealing the underlying pathological mechanisms of TED [17-20]. Using mass spectrometry profiling, differences in proteins between TED patients and control subjects can be investigated, potentially serving as biomarkers for aiding the detection and prognosis of TED in the future [21-24]. Furthermore, microbiomics research is indeed crucial as it sheds light on the connection between the gut microbiota and systemic immune responses that may influence the pathogenesis of TED, broadening the understanding of TED beyond the traditional central dogma and opening up new avenues for diagnosis, treatment, and potentially even prevention [25-27].

The emergence of multi-omics technologies offers a new perspective for studying TED, providing a more holistic and integrated view of the disease's pathophysiology. This comprehensive review aims to summarize the current knowledge regarding the application of multi-omics to TED, highlighting the potential for novel biomarker discovery and improved diagnostics.

# Methods

This systematic review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 Statement [28]. The review protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO) platform (registration number CRD42024533793). The protocol is available online at [https://www.crd.york.ac.uk/prospero/display\_rec ord.php?ID=CRD42024533793].

# Search Strategy

Two independent researchers (HYZ and YYZ) systematically searched the PubMed, Embase, and Web of Science databases from inception to February 10, 2024. All English-language publications were retrieved without restrictions on the country of origin or article type. The search strategy is provided in Supplementary Tables 1–3.

## **Selection Criteria**

We included only original studies, such as cross-sectional, retrospective, and prospective studies. The inclusion criteria were as follows: (a) studies involving patients with a confirmed diagnosis of TED, (b) studies employing any method of sequencing, including epigenomics, genomics, transcriptomics, microbiomics, metabolomics, and proteomics, and (c) studies comparing TED patients with a healthy control (HC) group and/or different subgroups within the TED population, such as those in the active TED phase versus the inactive TED phase. Studies were excluded if they met any of the following conditions: (a) review articles, animal studies, conference abstracts, meeting reports, studies with unclear information, and registry data, (b) studies focusing on diseases other than TED, (c) studies lacking explicit diagnostic criteria for TED, and (d) studies without sequencing results. Two researchers (HYZ and YYZ) independently selected the studies for exclusion. Conflicts were resolved through discussion to reach a consensus or by third-party arbitration.

## **Data Extraction and Visualization**

Data extraction was carried out independently by two researchers and subsequently double-checked. Disputes were resolved through discussion or by consulting a third specialist. Critical characteristics from all included studies were extracted and recorded. The extracted data encompassed the author, year of publication, region, sample type and size, age and sex of each group, sequencing methods, data analysis methods, and main findings.

#### **Risk of Bias and Quality Assessment**

The quality of the studies was assessed using the tool described by Lumbreras *et al.* according to QUADOMICS [29], which was developed to evaluate quality issues specific to omics research. Detailed information is presented in Supplementary Table 4, which considers sixteen criteria. The second and fourteenth items of QUADOMICS did not apply to the included studies, and none of the studies met item 12, which indicates that they interpreted the index test results with knowledge of the reference standard. Two independent authors (YYZ and YYD) conducted the quality assessment, and discrepancies were discussed until a consensus was reached. If consensus could not be achieved, a third specialist was consulted.

### Results

#### **Descriptive Characteristics**

A total of 412 studies were screened using the

described search strategy. At the end of the selection process, 69 studies were included in the systematic review. Details of the screening process are provided in the PRISMA flow diagram (Figure 1). Of the 69 studies, 4 studies (5.80%) utilized genomics and epigenomics, 33 studies (47.83%) utilized transcriptomics, 5 studies (7.25%) utilized metabolomics and lipidomics, 13 studies (18.84%) utilized proteomics, 7 studies (10.14%) utilized microbiomics, 6 studies (8.69%) utilized two omics assays, and 1 study (1.45%) utilized three omics assays. These studies were conducted between 2005 and 2024, with significant contributions from China, the UK, the US, Germany, Korea, and other countries. According to the quality assessment, 8 of the 69 studies received a score of "low" 54 received a score of "moderate" and 7 received a score of "high" Further information about this quality assessment is available in Supplementary Table 4.



Figure 1. Flow Diagram of study selection – adapted from the flow diagram template provided by PRISMA 2020.

 Table 1. Detailed information of studies on genomics and epigenomics

Author	Year	Region	Sample	Total	TED			Control			Sequencing	Data analysis	Main findings	
			types	sample size	Sample size	Gender (M/F)	Age (y)	Sample size	Gender (M/F)	Age (y)	methods	methods		
G Dottore et al.[30]	2021	Pisa, Italy	Orbital tissue (OF)	12	6	4/2	50.30 ± 12.50	6	4/2	65.20 ± 17.10	WGS	/	No gene variants were detected but the differentially expressed genes in TED are involved in the regulation of proliferation, apoptosis, and cell cycle, with an increase in global DNA methylation levels.	
S Virakul	2021	Rotterdam, Netherlands	Orbital tissue (OF)	12 F)	4 (Active)	1/3	63.50 ± 20.38	4	/	/	WGBS	Enrichment analysis of network, pathway and function	Active TED hypermethylated genes clearly linked to inflammation, while hypomethylated genes linked to adipogenesis and autoimmune-related genes.	
et al.[21]					4 (Inactive)	/	/							
Y Liang et al.[31]	2021	Shanghai, China	Orbital tissue (Connective tissue)	8	4	1/3	35.25 ± 6.34	4	0/4	39.00 ± 7.85	WGBS	KEGG, GO, Co-analysis of genes in methylation and expression level	In those differentially methylated genes, PTPRU and VCAM-1 performed significant alteration in methylation level and potentially participated in pathogenesis.	
Z Xin et al.[33]	2019	Beijing, China	Peripheral blood (PBMC)	12	6	0/6	54.5 (47- 61)	6	0/6	56.2 (46- 54)	RRBS	Enrichment analysis of pathway, methylation-based inference of regulatory activity score calculation, random forest analysis	Four significant pathways (Toxoplasmosis, Axon guidance, Focal adhesion, and Proteoglycans in cancer) were identified, with genes like LDLR, CDK5, and PIK3CB correlated with TED phenotype.	
Z Xin et al.[32]	2017	Beijing, China	Peripheral blood (Plasma)	12	6	6/0	55.00 ± 3.87	6	6/0	51.00 ± 2.58	RRBS	GO, KEGG, PCA	Several genomic loci were identified to have significant differences in methylation patterns that were associated with TED incidence, such as CD14, IL17RE, CDK5, DRD4, and ZCCHC6.	

OF: orbital fibroblast; EOM: extraocular muscle; PBMC: peripheral blood mononuclear cell; WGS: whole-genome sequencing; WGBS: whole-genome bisulfite sequencing; RNA-seq: RNA sequencing; RRBS: reduced representation bisulfite sequencing; KEGG: Kyoto encyclopedia of genes and genomes; GO: gene ontology; PCA: principal component analysis; TED: thyroid eye disease; GD: Graves' disease.

A total of 69 studies involved 1,363 TED patients and 1,504 controls (including 1,007 healthy controls (HCs) and 497 patients with GD or other diseases). The patients included in this review exhibit low heterogeneity, with ages ranging from 30.0 to 70.3 years and an overall mean age of approximately 48.40 years. The distribution of sex was similar across most studies, with a total of 322 males and 776 females. Although most of the included studies have a higher number of female participants, 2 studies reported a significantly higher proportion of male patients (male/female ratio  $\geq$  1.5). This data aligns with population statistics for TED.

The included studies analyzed various subgroups of TED. Forty studies compared TED with HC. Nine studies compared TED, GD, and HC. Five studies compared TED and GD. Six studies compared active TED, inactive TED, and HC. Seven studies examined TED based on severity, smoking status, which is a risk factor for the development of TED, and other valuable factors. One study compared TED, nonspecific orbital inflammation, sarcoidosis, and granulomatosis with polyangiitis. Additionally, one study employed 16S rRNA gene sequencing and metabolic network-driven analysis to investigate the association between gut microbiota and TED-related traits.

#### **Genomics and Epigenomics**

This systematic review included 5 studies

employing genomics and epigenomics. Of these, 1 study (20.0%) utilized genomics, while 4 studies (80.0%) investigated DNA methylation. Among the studies, 3 (60.0%) used orbital tissues as samples, and 2 studies (40.0%) analyzed peripheral blood.

In the genomic and epigenomic analysis of TED, whole-genome sequencing (WGS) and DNA methylation assays were applied, yet no significant genetic variants were identified [30]. For DNA methylation analysis, the sequencing approaches included whole-genome bisulfite sequencing (WGBS) [21], methylation microarrays [31], and reduced representation bisulfite sequencing (RRBS) [32, 33], DNA methylation data were analyzed using various approaches, including differential expression analysis to identify key genes [21], functional and pathway analysis with Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) for gene role classification [21, 31], network analysis to explore gene interactions [21], and epigenetic analysis techniques such as differentially methylated probes and differentially methylated region distribution to pinpoint methylation changes [31-33]. Bioinformatics tools like Ingenuity were used for data integration, while statistical and machine learning methods, including MIRA score calculation and random forest analysis, were employed for pattern recognition and correlation analysis [33], DNA methylation analysis detected various biomarkers. Notably, genes related to inflammation, adipogenesis, and autoimmunity,

such as PTPRU and VCAM-1, exhibited significant alterations in methylation levels, potentially playing a key role in the pathogenesis of TED [21, 31]. Pathway analysis and MIRA scoring identified four biological pathways significantly associated with TED, with genes such as LDLR, CDK5, and PIK3CB correlating with the TED phenotype [33]. Additionally, several genomic loci, including CD14, IL17RE, CDK5, DRD4, and ZCCHC6, exhibited significant differences in methylation patterns associated with TED incidence [32].

# Transcriptomics

This systematic review included 36 studies employing transcriptomics. Of these, 29 studies (77.8%) analyzed orbital tissues (orbital connective tissues in 27 studies, 75.0%, and extraocular muscles in 2 studies, 5.6%). Six studies (16.7%) used peripheral blood as the sample (peripheral blood mononuclear cells (PBMC) in 4 studies, 11.1%, and serum in 2 studies, 5.6%), and 1 study (2.8%) used thyroid tissue. Regarding sequencing methods, 16 studies (44.4%) utilized RNA-seq, 13 studies (36.1%) employed microarray, 4 studies (11.1%) used single-cell RNA sequencing (scRNA-seq), and 1 study (2.8%) implemented miRNA microarray. Additionally, nanostring and single-nucleus RNA sequencing (snRNA-seq) were each used in 1 study (2.8%). A wide range of data analysis methodologies were employed. DEGs were used in 31 studies (86.1%), KEGG pathway analysis in 20 studies (55.6%), GO analysis in 15 studies (41.7%), gene set enrichment analysis (GSEA) in 7 studies (19.4%), and protein-protein interaction (PPI) networks in 5 studies (13.9%). Cell-cell interaction analysis and principal component analysis (PCA) were each used in 2 studies (5.6%). Other methods, such as miRanda, competing endogenous RNA (ceRNA) network construction, and co-expression analysis, were employed in 1 study each (2.8%).

Most studies utilized a single approach– identifying DEGs in TED patients compared to HCs and conducting functional analysis using GO, KEGG, and GSEA. In studies analyzing orbital tissues, genes involved in antigen presentation, immune adhesion, interferon- $\gamma$  (IFN- $\gamma$ ) signaling, and immune cell markers for macrophages, B cells, and T cells were found to be upregulated. Additionally, the relaxin signaling pathway, which regulates fibrosis in TED, was active in TED patients. Four studies reported dysregulation of Wnt signaling gene expression, including Wnt5a, secreted frizzled-related proteins, and Dickkopf-related proteins [19, 34–36]. It is well established that the typical Wnt signal regulates the fate of neural crest progenitor cells. Similarly, activation of the Wnt/ $\beta$ -catenin pathway inhibits commitment to the adipocyte lineage while promoting differentiation into bone cells [37, 38]. Two studies used scRNA-seq to analyze peripheral blood mononuclear cells (PBMCs) and found that CD169+ classical monocytes and a novel CD4+ cytotoxic T lymphocyte (CTL) subtype with chemotactic and inflammatory characteristics contributed to hyperinflammation, fibrosis, and adipogenesis in orbital tissues [39, 40]. Additionally, Z Cheng *et al.* identified the VEGF-A gene as a regulator of the cytotoxic function of CD4+ CTLs in TED [41].

# Proteomics

This systematic review included 18 studies employing proteomics. Among these, 5 studies (27.8%) analyzed orbital tissues (orbital connective tissues in 4 studies, 80.0%, and extraocular muscles in 1 study, 20.0%). Four studies (22.2%) analyzed serum, and 9 studies (50.0%) used tear samples. Liquid chromatography-tandem mass spectrometry (LC-MS/MS) was the most used sequencing assay, employed in 11 studies (61.1%). Additionally, 2 studies (11.1%) utilized matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS), while other studies (27.8%) used Nano LC/QTOF, surface-enhanced laser desorption/ ionization time-of-flight mass spectrometry (SELDI-TOF-MS), and proximity extension assays.

Of the five studies that analyzed orbital tissues, four employed LC-MS/MS to examine differentially expressed proteins (DEPs) by comparing TED patients to HCs. These studies found that proteins involved in tissue inflammation, adipose tissue differentiation, and metabolism were typically overexpressed in TED patients [21, 23]. These findings suggest a shift in glycometabolism and lipometabolism in TED orbital tissues, which may help elucidate the underlying pathological mechanisms of TED. In serum samples, DEPs analysis, along with KEGG, GO, and PPI analyses, revealed significantly increased levels of IL6, CSF1, FLT3LG, and C4A in TED patients. These inflammatory proteins may play a crucial role in the pathogenesis of TED and could serve as potential new biomarkers for clinical use [42, 43]. Nine studies analyzed DEPs in tear samples, finding that protective proteins such as PROL1, PRP4, and β2-microglobulin were markedly downregulated, while inflammatory proteins such as lysozyme C and cystatin S were upregulated in the patient group compared to controls [44-47]. Enrichment analysis of the DEPs indicated that pathways related to the immune system, apoptosis, cell cycle, carbohydrate metabolism, and protein synthesis and degradation may be key in TED patients.

### Table 2. Detailed information of studies on transcriptomics

Author	Year	Region	Sample	Total	TED			Control	ontrol		Sequencin	Data analysis	Main findings
			types	sample size	Sample size	Gend er (M/F)	Age (y)	Sample size	Gend er (M/F)	Age (y)	g methods	methods	
L Zhu et al.[63]	2020	Beijing, China	Orbital tissue (EOM)	17	7	2/5	/	10	/	/	RNA-seq	GO, KEGG	The total m6A content was significantly increased in TED patients and identified aberrant expression of m6A methylation regulators.
X Bai et al.[35]	2022	Changsha, China	Orbital tissue (Connect ive tissue)	12	6	0/6	36.00 ± 13.45	6	3/3	30.67 ± 9.22	RNA-seq	GO, KEGG	Wnt was dysregulated and C5 was elevated in inactive TED vs HC, speculating C5 may promote remodeling of the orbital tissue.
A Sun et al.[64]	2023	Guangzho u, China	Orbital tissue (Connect ive tissue)	18	12	4/8	41.50 ± 13.50	6	3/3	44.50 ± 12.80	RNA-seq	KEGG, PPI	Relaxin signaling pathway is an important regulatory pathway in TED fibrosis pathogenesis, probably regulated by the differential expression of COLIA1, MMP2, FOS, GNG4, and CREB5.
H Ye et al.[65]	2022	Guangzho u. China	Orbital tissue	12	6 (Type I)	1/5	31.50 ± 7.60	/	/	/	RNA-seq	GO, KEGG, ceRNA	Hsa_circ_0007006 may regulate major genes, such as MMP2 and COL1A1.
[00]		u) enna	(Connect ive tissue)		6 (Type II)	3/3	51.50 ± 10.10					network construction, PPI	through the relaxin signaling pathway to affect TED typing.
L Wu et al.[66]	2021	Shanghai, China	Orbital tissue (Connect ive tissue)	12	6	/	/	6	/	/	RNA-seq	GO, KEGG, multivariate analysis of transcript splicing	Differential expression transcript and differential alternative splicing genes were found in TED vs HC, associated with immune modulation, extracellular matrix remodeling and adipogenesis.
J Khong et al.[67]	2015	Victoria, Australia	Orbital tissue (Connect ive tissue)	54	12 (Active) 21(Inact ive)	/	/	21	/	/	Microarray	GO, KEGG, GSEA	TIMD4, defensins (DEFA1, DEFA1B, and DEFA3) and markers for adipogenesis (SCD, FAD51, and SCDP1) were overexpressed in active TED vs inactive TED, associated with immune and inflammatory response.
L Wu et al.[68]	2020	Shanghai, China	Orbital tissue (Connect ive tissue)	6	3	/	/	3	/	/	RNA-seq	GO, KEGG, Co-expressio n and interaction analysis of circRNA-miR NA	The differentially expressed circRNAs might participate in pathogenesis of TED, and circRNA_14940-CCND1-Wnt signaling pathway might be an important regulatory axis.
Y Wang et al.[69]	2023	Wuxi, China	Orbital tissue (Connect ive tissue)	6	3	/	/	3	/	/	RNA-seq	PPI, GO, KEGG, GSEA	MAB21L1, PIK3C2G and CLVS2 were downregulated in TED vs HC, associated with dysregulation of differentiation, oxidative stress and developmental pathways.
L Wu et al.[70]	2021	Shanghai, China	Orbital tissue (Connect ive tissue)	6	3	/	/	3	/	/	RNA-seq	GO, KEGG, Co-expressio n analysis of lncRNA-mR NA	LncRNAs might regulate extracellular matrix remodeling in orbital tissue of TED. The target genes of IncRNAs involved in the lipid metabolism and cytokine-cytokine receptor interaction were identified. The negative correlations between Inc-PTP4A2-3:7-TNXB and LINC01139:4-SPON1 might be related to the altered extracellular matrix.
L Wu et al.[17]	2020	Shanghai, China	Orbital tissue (EOM)	10	5	/	/	5	/	/	Microarray	GO, KEGG	The expressions of genes involved in cell adhesion and ECM metabolism were significantly different between TED and HC.
M Chen et al.[18]	2008	Taipei, China	Orbital tissue (Connect ive tissue)	25	17	2/15	31.30 ± 6.20	8	2/6	41.00 ± 10.80	Microarray	GSEA	Lysosome-related genes, such as CLN2, CLN3 and HEXB, were downregulated in TED, may be involved in the pathogenesis of adipose tissue hypertrophy in TED.
P Mou et al.[71]	2018	Shanghai, China	Orbital tissue (Connect ive tissue)	15	10	/	/	5	/	/	RNA-seq	GO	PTX3 in orbital tissue and plasma was increased in TED vs HC, might playing a role in the pathogenesis of TED at least in the deposition of extracellular matrix.
W Tao et al.[34]	2017	Florida, US	Orbital tissue (Connect ive tissue)	9	5	/	38.22 ± 11.78	4	/	38.43 ± 9.16	RNA-seq	GO	In TED OASC, expression of early neural crest progenitor marker (Wnt signaling, ZIC genes and MSX2) was lost. Meanwhile, ectopic expression of HOXB2 and HOXB3 was found in the OASC from TED.
Z Li et al.[72]	2022	Guangzho u, China	Orbital tissue (Connect ive tissue)	20	10	/	/	10	/	/	scRNA-seq	Trajectory analysis, cell-cell interactions analysis, GO, KEGG	The study revealed the pro-inflammation and pro-adipogenesis role of LPFs with RASD1 expression. ACKR1* ECs and ATMs with an M2 phenotype were also identified as playing a role in

# Int. J. Biol. Sci. 2024, Vol. 20

Author	Voor	Region	Sample	Total	al TED Control Se					Sequencin Data analy	Data analyzaia	ysis Main findings	
Author	Iear	Region	types	sample size	Sample size	Gend er (M/F)	Age (y)	Sample size	Gend er (M/F)	Age (y)	g methods	methods	wan mungs
						(			(				TED. CD8+CD57+ CTLs showed the terminal differentiation phenotype and high IFN-g gene expression.
B Lee <i>et</i> <i>al.</i> [73]	2018	California, US	Orbital tissue (Connect ive tissue)	6 (Active)	3	0/3	70.33 ± 7.93	3	0/3	66.00 ± 9.93	RNA-seq	GO, KEGG, enrichment analysis of drugs and diseases	The IL-5 and chemokine signaling pathways were highly enriched, and very-low-density-lipoprotein receptor activity and statin medications were implicated as having a potential role in TED.
D Ezra et al.[19]	2012	London, UK	Orbital tissue (Connect ive tissue)	9 (Active)	5	1/4	51.20 ± 12.37	4	/	/	Microarray	/	The highest-ranked differentially expressed genes were dominated by IGF-1 signaling genes and data also demonstrate dysregulation of Wnt signaling gene expression, including Wnt5a, SFRPs and DKK.
D Kim <i>et</i> <i>al.</i> [36]	2021	Maryland, US	Orbital tissue (Intracon al connecti ve tissue)	6	3	1/2	46.33 ± 4.04	3	/	/	RNA-seq, single-nucle us RNA sequencing	KEGG, RNA velocity analysis, pseudotime analysis, regulon analysis, PCA	Signaling pathways including PI3K-Akt signaling, etc. are enriched in orbital fat isolated from patients with TED. The adipocyte-specific genes FABP4/5, APOE, PPARG and ADIPOQ during adipogenic differentiation differential expressed. The insulin-like growth factor-1 receptor and Wnt signaling pathways appear to be enriched early in adipogenesis.
T Planck et al.[20]	2014	Malmö, Sweden	Orbital tissue (Intraorb ital	26	8 (TED smoker s)	4/4	57.50 ± 9.35	5 (Healthy smokers)	0/5	/	Microarray	Enrichment analysis of pathway	Immediate early genes, IL-1B and IL-6 were overexpressed in smokers with severe active TED compared to nonsmokers, suggesting that smoking
			connecti ve tissue)		8 (TED nonsmo kers)	1/7	60.50 ± 14.07	5 (Healthy nonsmokers )	0/5	/			activates pathways associated with adipogenesis and inflammation.
T Planck et al.[74]	2011	Malmö, Sweden	Orbital tissue	30	10 (Active)	4/6	55.30 ± 11.25	10	1/9	52.90 ± 9.55	Microarray	Enrichment analysis of	PTHLH, which can inhibit adipogenesis, is downregulated both
			(Intraorb ital connecti ve tissue)		10 (Chroni c)	0/10	54.40 ± 12.55	10 (Chronic lymphedem a)	1/9	52.90 ± 9.55	,	pathway	in active and chronic TED, indicating the possibility of an increased risk of adipogenesis.
M Lantz et al.[75]	2005	Mainz, Germany	Orbital tissue (Intraorb ital connecti ve tissue)	10	5	2/3	54.60 ± 10.16	5	/	/	Microarray	/	Adipocyte-related immediate early genes are overexpressed in active ophthalmopathy, and CVR61 may have a role in both orbital inflammation and adipogenesis and serve as a marker of disease activity.
J Rosenbau m <i>et</i>	2015	Portland, US	Orbital tissue (Connect	86	25	/	51.62 (Mean)	20	/	61.00 (Mean)	Microarray	GSEA, PCA	Inflammatory markers are far less characteristic of TED relative to other orbital inflammatory diseases
al.[76]			ive tissue)					25 (NSOI)	/	50.74 (Mean) 41.7			oronal inflatinitatory diseases.
								8	/	(Mean)			
								(Sarcoidosis )	/	(Mean)			
R Ma et al.[77]	2022	Shanghai, China	Orbital tissue (Connect ive tissue)	10	5	/	/	5	/	/	RNA-seq	KEGG	Multiomic analyses of glycolysis and mitochondrial metabolism (TCA & OXPHOS) suggested that a glycolytic shift may exist in TED.
S Jang et al.[78]	2016	Seoul, Korea	Orbital tissue (Connect ive tissue)	12	8	/	/	4	/	/	Microarray	/	The level of miR-146a expression was significantly higher in TED orbital adipose tissue than in non-TED.
S Kumar et al.[79]	2005	Minnesota, US	Orbital tissue (Connect ive tissue)	28	20	/	/	8	/	/	Microarray	/	Gene expression profiling identified sFRP-1 as another gene up-regulated in TED patient tissues.
J Kor et al.[80]	2023	Seoul, Korea	Orbital tissue (Connect ive tissue)	18	12	5/7	21-59	6	2/4	27-64	RNA-seq	GO, KEGG, GSEA	YAP expression in TED was enhanced. Response to mechanical stimulus-related genes were overexpressed in TED, along with those enriched for adipose proliferation, inflammatory responses and hormone stimulus responses.
S Fang et al.[8]	2019	Shanghai, China	Orbital tissue (Connect ive	6	3	/	/	3	/	/	scRNA-seq	/	Gene expression of the master transcription factors TBX21 and STAT1 (Th1 cell), GATA3 and STAT6 (Th2 cell), RORA and STAT3 (Th17

https://www.ijbs.com

Author	Year	Region	Sample	Total sample size	TED			Control			Sequencin Data analy		ysis Main findings
		8	types		Sample size	Gend er (M/F)	Age (y)	Sample size	Gend er (M/F)	Age (y)	g methods	methods	
			tissue)			(14,1)			(1441)				cell), and AHR (Th22 cell) revealed the possible involvement of various effector T cells in TED orbits.
Z Yue et al.[81]	2022	Shanghai, China	Orbital tissue (Connect ive tissue)	6	3	/	/	3	/	/	RNA-seq	GO, KEGG, target gene prediction	A total of 50 tRFs and 361 mRNAs were dysregulated in the TED group, and tRF5-GluCTC, PMAIP1, HSD17B2 and ATF3 were verified to be significantly differentially expressed in TED.
Z Yue et al.[82]	2021	Shanghai, China	Orbital tissue (Connect ive tissue)	6	3	/	/	3	/	/	Microarray	PPI, GO, KEGG, ceRNA network construction	Two important ceRNA pathways were identified, the LINC01820:13-hsa-miR-27b-3p-FPR2 ceRNA pathway and the ENST00000499452-hsa-miR-27a-3p-CX CL1 pathway, which probably affect the autoimmune response and inflammation in TED patients.
D Wang et al.[39]	2024	Guangzho u, China	Peripher al blood (PBMC)	9	5	/	/	4 (GD)	/	/	scRNA-seq	KEGG, pseudotime trajectory analysis, transcription factor analysis, cell-cell interactions analysis,	Monocytes along with CD169 <sup>+</sup> clas_mono-derived macrophages, contributed to hyperinflammation, fibrosis, and adipogenesis in orbital tissues. CD169 <sup>+</sup> clas_mono expansion is related to activated IFNg signaling pathway in TED.
N Kim et al.[83]	2023	Korea (Seongnam , Seoul)	Peripher al blood (Plasma)	15	5 (Active) 5 (Inactiv e)	3/2 3/2	$46.40 \pm 11.20$ $43.00 \pm 16.60$	5	3/2	39.60 ± 12.20	NanoStrin g	KEGG	Ten circulating miRNAs were found to be differentially expressed, with miR-484 and miR-192-5p showing significantly lower expression levels in the active TED group compared to the inactive TED group.
L Zhang et al.[59]	2018	UK (Cardiff), Italy (Milan), Germany (Essen)	Peripher al blood (Plasma)	46	19	19/0	48.40 ± 10.60	13 14 (GD)	13/0 14/0	47.2 ± 11.7 51.6 ± 17.9	miRNA-se q	Lasso regression, KEGG, target gene prediction, PCA	The top known and novel miRNAs were hsa-mir-497, hsa-mir-320b-1 and hsa-mir-320b-2, up-regulated in GD, Novel:19_15038 and hsa-miR-27a-3p up-regulated in TED.
R Liu et al.[84]	2023	Shanghai, China	Peripher al blood (PBMC)	20	10	/	/	10	/	/	Microarray	Target gene prediction	MiR-376b, which is downregulated in TED patients' PBMCs and regulated by T3, might play a role in TED pathogenesis by influencing HAS2 and inflammatory factors.
Y Wang et al.[40]	2021	Xi'an, China	Peripher al blood (PBMC)	9	6	/	/	3 (GD)	/	/	scRNA-seq	GSEA, GO	The novel TED-specific cell type CD4 <sup>+</sup> cytotoxic T lymphocytes, which are characterized by chemotactic and inflammatory features, were characterized by cytotoxicity (GZMB, KLRG1, GZMH, PRF1, GNLY and FGFBP2), chemotaxis (CCL4, CCL5 and CX3CR1) and inflammation (IL-1B and IFNG).
Z Cheng et al.[41]	2023	Xi'an, China	Peripher al blood (PBMC)	40	20	/	/	20 (GD)	/	/	RNA-seq	PPI, GSEA	VEGC-A was identified as a regulator of the cytotoxic function of CD4 <sup>+</sup> cytotoxic T lymphocytes in TED.
L Wescomb e <i>et al.</i> [85]	2009	Sydney, Australia	Thyroid	18	10	1/9	40.6 (20– 57)	8 (GD)	0/8	28 (17-44)	Microarray	/	Two hundred and ninety-five genes were differentially expressed between patients with TED and GD. Of these, the cardiac calsequestrin gene was the most highly expressed gene in TED.
Y Huang et al.[9]	2022	Shanghai, China	Orbital tissue (Lacrima l Gland)	76	26 (Active) 25 (Inactiv e)	13/13 9/16	48.92 ± 11. 59 41.84 ± 12. 54	25	12/13	42.12±11. 66	RNA-seq	GO, KEGG	Increased expression of cytokines and chemokines accompanied by a variety of immune cell infiltrations mainly involving T cells, B cells, and monocytes was found in TED lacrimal glands. An in-depth investigation into T cell subsets revealed IFN-γ-producing Th1 and IL-17A-producing Th17 cell-dominated autoimmunity in the active TED lacrimal microenvironment.

EOM: extraocular muscle; PBMC: peripheral blood mononuclear cell; PPI: protein-protein interaction network; GSEA: gene set enrichment analysis; scRNA-seq: single cell RNA sequencing; miRNA-seq: microRNA sequencing; NSOI: nonspecific orbital inflammation; GPA: granulomatosis with polyangiitis; TED: thyroid eye disease; GD: Graves' disease; GO: Gene Ontology.

Author	Year	Region	Sample	Total	Total TED Conduct Act C				Control			Data	Main findings
			types	sampl e size	Sample size	Gender (M/F)	Age (y)	Sample size	Gender (M/F)	Age (y)	methods	analysis methods	
S Virakul et al.[21]	2021	Rotterdam, Netherlands	Orbital tissue (OF)	14	4 (Active)	1/3	63.50 ± 20.38	5	0/5	61.20 ± 17.61	LC-MS/MS	Enrichment analysis of	Proteins that are typically involved in inflammation,
					5 (Inactive)	0/5	48.20 ± 12.12					network, pathway and function	cellular proliteration, hyaluronan synthesis and adipogenesis, were overexpressed, while various proteins associated with extracellular matrix biology and fibrotic disease, were typically overexpressed.
L Wu et al.[17]	2020	Shanghai, China	Orbital tissue (EOM)	6	3	/	/	3	/	/	LC-MS/MS	GO, KEGG, GSEA, PPI	Proteins mainly related to the composition (such as MYH1) and contraction force (such as MYH1) of the muscle fibers were significantly up regulated in TED, as well as those (such as VCAN) associated with cell adhesion. Differentially expressed proteins related to the components and metabolism of ECM (such as COL1A1) were identified in EOM samples of TED.
K Cheng et al.[23]	2013	Kaohsiung, China	Orbital tissue (Connectiv e tissue)	14	7	4/3	56.60 ± 9.70	7	2/5	57.60 ± 7.10	LC-MS/MS	/	Up-regulations of the specific proteins are associated with biochemical mechanisms or capacities against endoplasmic reticulum stress, mitochondria dysfunction, and cell proliferation as well as apoptosis in TED orbital fat tissues.
N Matheis <i>et al.</i> [86]	2015	Sweden (Malmö), Germany (Rhinel-Palati	Orbital tissue (Connectiv e tissue)	64	6 (Untreated nonsmokers)	1/5	30 (20 -73)	12 (Control orbital tissue)	4/8	67.2 (48–85)	MALDI TOF/TOF	/	Proteins involved in tissue inflammation, adipose tissue differentiation, lipid metabolism, and tissue remodeling were up
		nate)			16 (Steroid-treated nonsmokers)	5/11	55.1 (20-76)	13 (Control peripheral	1/12	61 (16-78)			untreated patients with TED and steroids decreased the expression of these proteins, whereas
					17 (Steroid-treated smokers)	7/10	48.5 (27-71)	tissue)					smoking attenuated such effect.
R Ma et al.[77]	2022	Shanghai, China	Orbital tissue (Connectiv e tissue)	10	5	/	/	5	/	/	LC-MS/MS	KEGG	Multiomic analyses of glycolysis and mitochondrial metabolism (TCA & OXPHOS) suggested that a glycolytic shift may exist in TED.
J Kang et al.[42]	2021	Kunming, China	Peripheral blood (Plasma)	9	3	2/1	42–54	3 3 (GD)	0/3 1/2	25–27 34–50	LC-MS/MS	GO, KEGG, PPI	MYH11, P4HB, and C4A were markedly upregulated in TED patients and have been reported to participate in apoptosis, autophagy, the inflammatory response, and the immune system.
L Zhang et al.[59]	2018	UK (Cardiff), Italy (Milan),	Peripheral blood	46	19	19/0	48.40 ± 10.60	13	13/0	47.2 ± 11.7	Nano LC/QTOF	Lasso regression,	Proteomic and miRNA analyses identified circulating biomarkers,
		Germany (Essen)	(Plasma)					14 (GD)	14/0	51.6± 17.9		KÉGG	including Zonulin and Fibronectin, differentiating between GD, TED, and healthy controls, suggesting diagnostic and prognostic potential.
H Ueland et al.[43]	2022	Bergen, Norway	Peripheral blood	220	36	7/29	41 (19- 70)	120	54/66	40 (23- 68)	Proximity extension	Student's t-test or	Distinctly increased levels of IL6, CSF1, FLT3LG, and FGF21 were
			(Plasma)				,	64 (GD)	16/48	43 (15- 66)	assay	Mann- Whitney U-test, Spearman's correlation, ROC	observed in TED, suggesting that these inflammatory proteins could be important in the pathogenesis, and therefore potential new biomarkers for clinical use.
T Shi et al.[61]	2023	Beijing, China	Peripheral blood (Plasma)	60	30	16/14	46.50 ± 10.60	30	15/15	42.10 ± 9.00	LC-MS/MS	Lasso regression, KEGG, interaction analysis of metabolites and gene	CPS1, GP1BA, and COL6A1 were identified as feature proteins and had better prediction performance for TED compared to the baseline model.
T Shi et al.[87]	2022	Beijing, China	Tear (Exosomes)	64	24	9/15	45.20 ± 10.10	16 24 (GD)	5/11 10/14	40.40 ± 9.60 40.30 ± 8.90	LC-MS/MS	GO, KEGG	The levels of the exosomal proteins Caspase-3, complement C4A and APOA-IV were significantly increased in TED
<u> </u>	2022	<u>cı</u> t.:		(0)	20	0./22	10.00	20	0./22	10.57	1010/00	VECC 25	patients compared to GD patients and controls.
X Zhou et	2022	Shanghai,	Tear	60	30	8/22	42.00 ±	30	8/22	40.27 ±	LC-MS/MS	KEGG, GO	Up-regulated proteins were

Author	Year	Region	Sample	Total	Fotal TED Co			Control			Sequencing	Data	Main findings
		C	types	sampl e size	Sample size	Gender (M/F)	Age (y)	Sample size	Gender (M/F)	Age (y)	methods	analysis methods	U U
al.[88]		China					11.91			9.32			mainly enriched in lipid metabolism and platelet molecular function, and down-regulated proteins were involved in binding, cell junction, and cellular process.
C Aass et al.[47]	2016	Oslo, Norway	Tear	42	21	6/15	44 (26– 69)	21 (GD)	4/17	57 (20– 77)	LC-MS/MS	Spearman's correlation	Upregulation of lacrimal gland proteins such as lysozyme C, lacritin, antileukoproteinase and zinc-alpha-2-glycoprotein 1 suggests involvement of the lacrimal gland in the pathogenesis of TED.
N Matheis <i>et al.</i> [44]	2012	Mainz, Germany	Tear	60	45	8/37	50 (22– 77)	15	2/13	45 (33– 74)	SELDI-TOF- MS, MALDI-TOF -MS	Analysis of variance, ROC	PRP4 and $\beta$ 2-microglobulin was decreased while lysozyme C and cystatin S was upregulated in TED, demonstrating altered regulation of proinflammatory and protective proteins in tears of patients with TED.
L Jiang et al.[89]	2021	Shanghai, China	Tear	12	6	3/3	50.20 ± 5.40	6	2/4	46.70 ± 6.30	LC-MS/MS	PLS-DA, GO, KEGG, PPI, GRN	Signaling pathways of the immune system, apoptosis, cell cycle, cytoskeleton, metabolism of carbohydrates, cofactors and vitamins, protein synthesis and degradation, vesicle-mediated transport of membrane and proteins may play key roles in patients with inactive TED.
C Chng et al.[90]	2018	Singapore, Singapore	Tear	72	18 (Mild)	5/13	54.17 (Mean)	18 (AITD)	5/13	57 (Mean)	LC-MS/MS, SWATH-MS	Mann-Whit	Two proteins, S100A4 and PIP showed consistent dysregulation
					18 (Severe)	11/7	56 (Mean)	18	8/10	58 (Mean)		ney U test, Kruskal-W allis tests, geometric mean	trends in the discovery and validation phase experiments, may serve as potential biomarkers to predict progression to severe TED in patients with AITD.
E Kishazi et al.[46]	2018	Lausanne, Switzerland	Tear	12	6	/	/	6	/	/	LC-MS/MS	/	Cystatin c (TED/controls ratio: 1.53), alpha-1 antichymotrypsin (ratio: 1.70) and retinal dehydrogenase (ratio: 0.68), displaying differential levels in the tears of TED patients as highly promising biomarkers after verification.
R. Okrojek et al.[91]	2009	Mainz, Germany	Tear	60	45	/	/	15	/	/	SELDI-TOF- MS	Multivariat e discrimina nt analysis, ROC	A set of biomarkers in the mass range of 3000–20000 Da, showing significant differences. The corresponding ROC curve had an area under the curve of 0.99 and a specificity and sensitivity of over 90% each.
N Matheis	2015	Mainz, Germany	Tear	ur 120 3 	30	7/23	45.5 (17-68)	30	3/27	47.5 (21-70)	MALDI-TOF MS	Enrichment analysis of	Upregulation of inflammatory proteins and downregulation of
et al.[45]					30 (TED + Dry eye)	5/25	51 (31- 70)	30 (Dry eye)	7/23	54.5 (32-80)		pathway	protective proteins (i.e., PROL1/PRP4).

OF: orbital fibroblast; EOM: extraocular muscle; MALDI TOF/TOF: matrix-assisted laser desorption/ionization time-of-flight/ time-of-flight; Nano LC/QTOF: nanoscale liquid chromatography/quadrupole-time-of-flight; SELDI-TOF-MS: surface-enhanced laser desorption/ionization time-of-flight mass spectrometry; MALDI-TOF MS: matrix-assisted laser desorption/ionization time-of-flight mass spectrometer; SWATH-MS: sequential window acquisition of all theoretical spectra mass spectrometer; ROC: receiver operating characteristic; TED: thyroid eye disease; GD: Graves' disease; GO: Gene Ontology.

#### **Metabolomics and Lipidomics**

This systematic review included 9 studies employing metabolomics. Among these, 3 studies (33.3%) used targeted metabolomics, while 6 studies (66.7%) utilized non-targeted metabolomics, including 1 study (11.1%) that specifically used non-targeted lipidomic metabolomics. Of the 9 studies, 3 (33.3%) analyzed orbital connective tissues, 3 (33.3%) used serum, 1 study (11.1%) analyzed feces, 1 study (11.1%) employed tear samples, and 1 study (11.1%) examined both orbital connective tissues and serum.

The non-targeted lipidomic metabolomics study utilized nanoflow ultrahigh pressure liquid chromatography-electrospray ionization tandem mass spectrometry (nUPLC-ESI MS/MS) sequencing, with data analyzed using Student's t-test and PCA. This study identified significantly increased levels of sphingosine-1-phosphate in serum and urine samples, indicating its potential as a biomarker for TED diagnosis. Among the other 6 non-targeted metabolomics studies, 3 studies (50.0%) employed LC-MS sequencing, 2 studies (33.3%) used gas chromatography-time-of-flight mass spectrometry (GC-TOF MS), and 1 study (16.7%) utilized nuclear magnetic resonance (NMR) spectroscopy. Metabolite profiles in TED patients, including fumarate, proline, phenylalanine, and glycerol, suggested a potential metabolic connection between orbital connective tissues and blood metabolites. Notably, cholesterol metabolism was significantly linked to TED pathogenesis.

Table 4. Detailed information of studies on metabolomics and lipid	omics
--	-------

Author	Year	Region	Sample	Total	al TED C		Control			Sequencing	Data analysis	Main findings	
		-	types	sample size	Sample size	Gender (M/F)	Age (y)	Sample size	Gender (M/F)	Age (y)	methods	methods	-
F Biscarini et al.[27]	2023	UK (Wales, Newcastle, London), Italy (Milan, Pisa), Belgium (Brussels), Germany (Essen)	Fecal microbiota	146	10	/	/	11 26 (GD)	/ /	/	_NMR	PLS-DA	Short-chain fatty acids (SCFAs, butyrate, propionate etc.) was increased in TED vs HC.
R Du et al.[22]	2023	Shanghai, China	Orbital tissue (Connective tissue)	12	7	2/5	33.80 ± 8.90	5	1/4	39.80 ± 8.90	LC-MS/MS	PLS-DA, enrichment analysis of pathway	In TED vs HC, the phosphoenolpyruvic acid was upregulated and downregulated metabolites were uric acid, phosphoenolpyruvic acid etc. and the TED-related pathways identified included purine metabolism, beta-alanine metabolism, glutathione metabolism.
J Huang et al.[24]	2022	Shanghai, China	Orbital tissue (Connective tissue)	23	11	3/8	54.63 ± 14.49	12	2/10	36.58 ± 13.05	LC-MS/MS	PLS-DA, KEGG, interaction analysis of metabolites and gene	TED has a significant difference in the metabolic profiles, altering cellular metabolome, especially cholesterol metabolism.
R Ma et al.[77]	2022	Shanghai, China	Orbital tissue (Connective tissue)	10	5	/	/	5	/	/	LC-MS/MS	/	Metabolic profiles provided evidence that the parameters favoring glycolysis (F6P/F16BP, AMP/ATP, ADP/ATP, and lactate) were upregulated in the TED.
D Ji et al.[58]	2018	Seoul, Korea	Orbital tissue (Connective tissue)	10	5	3/2	53.6 (Mean)	5	3/2	52.8 (Mean)	GC-TOF MS	/	The up-regulated metabolites included proline, fumarate, and phenylalanine that were consistent with the expression pattern in blood metabolite analysis. The activated tissue metabolism was represented by amino acids, including asparagine, valine, allo-threonine, methionine, and glycine.
D Ji et al.[58]	2018	Seoul, Korea	Peripheral blood (Plasma)	79	26	7/19	39.50 ± 10.30	32 21 (GD)	6/26 6/15	/ 36.40 ± 10.40	GC-TOF MS	PLS-DA, ROC, enrichment analysis of pathway, PCA	With combination of proline and 1,5-anhydroglucitol, which were identified as TED-specific modulators, the re-constructed biomarker model greatly improved the statistical power and facilitated simultaneous discrimination among healthy control, TED, and GD without TED groups.
T Shi et al.[61]	2023	Beijing, China	Peripheral blood (Plasma)	60	30	16/14	46.50 ± 10.60	30	15/15	42.10 ± 9.00	LC-MS/MS	PLS-DA, lasso regression, KEGG, interaction analysis of metabolites and gene, PCA	Lycine, glycerol 3-phosphate, and estrone sulfate were identified as feature metabolites and had better prediction performance for TED compared to the baseline model.
S Byeon et al.[92]	2022	Seoul, Korea	Peripheral blood (Plasma)	86	31	8/23	/	32 23(GD)	6/26 6/17	/	nUPLC-ESI MS/MS	PCA	The sphingosine-1-phosphate levels showed consistent and significant increases in both plasma and urine samples of patients with TED.
H Ueland et al.[93]	2023	Bergen, Norway	Peripheral blood (Plasma)	200	36	8/23	/	100 64 (GD)	23/77 6/17	39 (15-70) /	LC-MS/MS	Linear regression models, factor analysis	No difference in systemic activation of the kynurenine pathway in GD patients with and without TED implies that the local Th1 immune response in the orbit is not
B Billiet et al.[54]	2022	Angers, France	Tear	45	21 (Active) 24 (Inactive)	5/16	55 (Mean) 51 (Mean)	/	/	/	LC-MS/MS	/	retilected systemically. Two short chain acylcarnitines, propionylcarnitine and butyrylcarnitine, and spermine showed increased concentrations and the ratio putrescine/ornithine, representing the activity of ornithine decarboxylase, was significantly increased in patients with active compared to inactive TED.

NMR: nuclear magnetic resonance; LC-MS/MS: liquid chromatography-mass spectrometry/mass spectrometry; GC-TOF MS: gas chromatography-time of flight mass spectrometry; nUPLC-ESI MS/MS: nanoflow ultrahigh pressure liquid chromatography tandem mass spectrometry; PLS-DA: partial least squares-discriminant analysis; ROC: receiver operating characteristic; TED: thyroid eye disease; GD: Graves' disease.

When comparing TED patients to HCs, metabolites such as short-chain fatty acids, uric acid, uracil, hexose-phosphate, and D-sedoheptulose 1,7-bisphosphate were enriched in TED samples. Combining TED-specific modulators-proline and 1,5-anhydroglucitol-with key metabolites-lycine, glycerol 3-phosphate, and estrone sulfate substantially improved the biomarker model's ability to discriminate between HCs, GD, and TED groups. All 3 targeted metabolomics studies utilized LC-MS/MS sequencing, with data analyzed using factor analysis, linear regression models, and various univariate analyses. Metabolic profiling revealed glycolysis-related significant upregulation of parameters (F6P/F16BP, AMP/ATP, ADP/ATP, and lactate) in orbital fibroblasts of TED patients compared to HCs. Additionally, active TED patients showed an increased ratio of putrescine to ornithine and spermine in serum compared to inactive TED patients.

# **Microbiomics**

This systematic review included 8 studies that utilized microbiomics. Among these, 6 studies (75.0%) analyzed fecal microbiota, 1 study (12.5%) examined orbital adipose tissues, and 1 study (12.5%) investigated ocular microbiota. Of the studies, 6 (75.0%) employed 16S rRNA sequencing, while 2 studies (25.0%) used 16S rDNA sequencing. The methodologies for identifying biomarkers and analyzing co-occurrence patterns within the microbiota varied. Notably,  $\alpha$ - and  $\beta$ -diversity indices were used in 5 studies (62.5%), and random forest analysis was employed in 3 studies (37.5%). Additionally, specific techniques such as KEGG, weighted gene co-expression network analysis (WGCNA), and module-trait association analysis were utilized in 1 study (12.5%).

The collective evidence from these studies suggests that gut microbiota may play a significant role in TED, with specific microbial imbalances potentially contributing to immune dysregulation and disease development. TED patients exhibit decreased bacterial community diversity but show increased proportions of Actinobacteria, Bacillus, Brevundimonas, higher *Firmicutes*-to-*Bacteroidetes* and а ratio compared to HCs [48, 49]. Specific taxa, such as Deinococcus-Thermus, Chloroflexi, and various bacterial species, display differential abundance patterns between TED and GD, indicating disease-specific microbial signatures [50]. Further characterization reveals distinct microbiome types unique to TED patients, with taxa like Klebsiella pneumoniae, Paracoccus, and Hemophilus correlating with disease

severity [48, 51]. Additionally, s\_Prevotella\_copri and f\_Prevotellaceae show significant correlation with TRAb, suggesting their potential role in TED development [52]. Furthermore, bacterial species such as *Bacteroides*, *Dialister*, and *Lactobacillus* exhibit associations with serum lipopolysaccharide-binding protein concentration, indicating their involvement in TED pathophysiology [53].

# Discussion

TED is an autoimmune inflammatory disorder of the orbit, characterized by inflammation and a range of symptoms that can impact vision and appearance. Despite the presence of clinical scoring systems, there remains a critical need for more objective biomarkers to improve early diagnosis, disease monitoring, and therapeutic efficacy evaluation. Recent advancements in high-throughput sequencing technology and the rapid accumulation of omics data have led to increased utilization of genomics, proteomics, and other omics approaches to identify novel targets and biomarkers. Our systematic review comprehensively analyzed studies that employed genomics, epigenomics, transcriptomics, proteomics, metabolomics, lipidomics, and microbiomics to identify potential biomarkers in various biological samples, including orbital tissues, peripheral blood, feces, and tears. The integration of multi-omics data has significantly advanced our understanding of the molecular mechanisms underlying TED and has facilitated the identification of new diagnostic and prognostic biomarkers.

Among the 69 articles included in our study, numerous biomarkers were identified across various mediums, including fecal microbiota, tear fluid, orbital tissues, blood, and thyroid tissues. Connective tissues and extraocular muscles are the primary sites of pathological processes in TED. Studies often focus on exploring disease-specific molecular changes such inflammation, adipogenesis, and immune as responses. For instance, through transcriptomic analysis, researchers have identified the upregulation of genes related to antigen presentation, immune cell adhesion, and interferon-y signaling in TED patients [19, 36]. Tear samples reflect the health status of the ocular surface. Proteomic analysis has revealed the downregulation of protective proteins and upregulation of inflammatory proteins, which may be associated with ocular surface inflammation in TED patients [54]. Fecal samples are used to investigate the association between gut microbiota and TED, revealing specific microbial imbalances related to immune dysregulation and disease development [50].

#### Table 5. Detailed information of studies on microbiomics

Author	Year	Region	Sample	Total	tal TED C			Control			Sequencing	Data analysis	Main findings	
		U	types	sample size	Sample size	Gender (M/F)	Age (y)	Sample size	Gender (M/F)	Age (y)	methods	methods	U U	
T Shi et al.[49]	2019	Beijing, China	Fecal microbiota	65	33	17/16	46.00 ± 11.71	32	16/16	43.40 ± 9.70	16S rRNA	<ul> <li>α- and</li> <li>β-diversity</li> <li>indexes,</li> <li>random forest</li> <li>analysis,</li> <li>Mann-Whitney</li> <li>U test, LEfSe</li> </ul>	Bacterial community diversity was significantly reduced in TED vs HC; <i>Bacteroidetes</i> was significantly increased in TED vs HC.	
T Shi et al.[50]	2020	Beijing, China	Fecal microbiota	95	33	17/16	46.00 ± 11.70	32 30 (GD)	16/16	$43.40 \pm 9.70 \\ 45.00 \pm 12.80$	16S rRNA	α- and β-diversity indexes, random forest analysis, Mann-Whitney U test, KEGG	Deinococcus-Thermus and Chloroflexi were significantly reduced in TED vs GD; Subdoligranulum and Bilophila were increased in TED vs GD; Viral protein family was the only enrichment metabolic pathway in GD vs TED.	
T Shi <i>et</i> al.[52]	2019	Beijing, China	Fecal microbiota	31	31	15/16	45.40 ± 11.80	/	/	/	16S rRNA	WGCNA, pearson correlation analysis, negative binomial regression, latent class analysis	The abundance of s_Prevotella_copri and f_Prevotellaceae was positively correlated with TRAb level.	
Q Zhang et al.[51]	2023	Changsha, China	Fecal microbiota	80	20 (Mild)	7/13	37.50 ± 10.05	18	3/15	33.50 ±	16S rDNA	α- and β-diversity	α- and β-diversity	No remarkable difference in diversity in TED vs HC; The
					25 (Moderate-to-severe) 17 (Sight threatoning)	8/17 12/5	40.00 ± 14.45 55.00 ±			10.60		indexes, PCA, KEGG, target gene prediction	abundance of Klebisiella_Pneumoniae was positively correlated with disease severity.	
F Biscarini et al.[27]	2023	UK (Wales, Newcastle, London), Italy (Milan, Pisa), Belgium (Brussels), Germany (Essen)	Fecal microbiota	146	46	6/40	47.04 ± 11.43	41 59 (GD)	9/32	46.08 ± 13.45 46.39	16S rRNA	α- and β-diversity indexes, random forest	Actinobacteria and F:B ratio were significantly increased and <i>Baceteroidetes</i> significantly decreased in GD/TED vs HC;	
								. ,		± 14.38		analysis, pearson correlation analysis	Bacteroides abundance was positively correlated with TSH and negatively correlated with free thyroxine; Presence of <i>Clostridiales</i> at diagnosis was correlated with long-term TRAb persistence post antithyroid treatment.	
A Fenneman	2023	Amsterdam, Netherlands	Fecal microbiota	15	8 (Low plasma levels of LBP)	2/6	42.12 ± 12.76	/	/	/	16S rRNA	Spearman correlation	Plasma LBP level was positively correlated with Gram-negative	
et al.[53]					7 (High plasma levels of LBP)	3/4	51.43 ± 9.31					analysis	gut bacteria abundance (Baceteroides spp., Dialister spp.); Lactobacillus spp. was strongly increased in H-LBP vs L-LBP.	
Y Li et al.[94]	2022	Beijing, China	Orbital tissue (Connective tissue)	54	27	9/18	48 (22-74)	27	17/10	25 (3- 67)	16S rRNA	a-diversity indexes, PCA, co-occurrence network, multiple linear regression and mixed effect model	Four microbiome types were identified in orbital fat, one of which was exclusive to TED patients. A higher proportion of <i>Pseudomonas</i> in TED, but a lower proportion of <i>Enhydrobacter</i> . <i>Flavobacterium</i> was positively correlated with chemosis.	
X Ji et al.[48]	2022	Taiyuan, China	Ocular microbiota	69	47	8/39	44.25 ± 13.44	22	7/15	62.64 ± 7.11	16S rDNA	α- and β-diversity indexes, spearman correlation analysis	The average relative abundance of <i>Bacillus</i> and <i>Brevundimonas</i> increased significantly in the TED group. <i>Corynebacterium</i> had a significantly decreased relative abundance. <i>Paracoccus,</i> <i>Hemophilus, Lactobacillus,</i> and <i>Bifidobacterium</i> were positively correlated with the severity of clinical manifestations or disease activity.	

LBP: lipopolysaccharide-binding protein; 16S rRNA: 16S ribosomal RNA; 16S rDNA: 16S ribosomal DNA; LEfSe: linear discriminant analysis effect size; WGCNA: weighted gene co-expression network analysis; GD: Graves' disease.

Through scRNA-seq, blood samples are utilized to explore the connections between the peripheral immune landscape and TED, such as the activation of CD169<sup>+</sup> classical monocytes, which may be pivotal in driving the autoimmune reactions characteristic of TED [39]. Combining data from different biological samples can provide a more comprehensive understanding of disease mechanisms. For example, combining orbital tissue and blood samples can reveal both local and systemic changes. Samples that are easily obtainable (such as blood and tears) may aid in developing non-invasive diagnostic tools for disease monitoring and prognosis. Examining biological samples from various sources allows for a more thorough comprehension of TED, spanning from localized tissue effects to broader immune system impacts, ultimately guiding the identification of more effective biomarkers.

In omics analyses, the selection of appropriate control groups is crucial for elucidating biological differences. In this systematic review, many studies compared TED patients with HC and patients with GD to identify biomarkers associated with TED. Comparing TED patients to HC helps reveal disease-specific molecular alterations, with the HC group providing a baseline against which pathological changes in TED patients can be contrasted. For example, significant differences in gene expression, protein levels, metabolites, and microbiota composition between TED patients and HC may shed light on pathogenic mechanisms and disease progression. Conversely, comparisons with GD patients aid in distinguishing the specific manifestations of TED within the context of GD. Although GD is a commonly associated with TED, not all GD patients develop TED. Differentiating TED from GD helps identify molecular signatures unique to TED, which may be related to disease severity, inflammatory responses, or tissue remodeling. Additionally, some studies compared TED with other inflammatory conditions, such as non-specific orbital inflammation, sarcoidosis, and granulomatosis with polyangiitis, or grouped patients based on factors like smoking status and serum lipopolysaccharidebinding protein levels. These comparisons aim to uncover the unique pathological features and influencing factors of TED.

In exploring biomarkers related to TED, the choice of sequencing methods is pivotal, requiring a nuanced understanding of different biological molecular levels as described by the central dogma, which is a principle that describes the flow of genetic information within a biological system [16]. The central dogma was initially proposed by Francis Crick in 1957 and has since been refined. It outlines the flow of genetic information from DNA to RNA to proteins, and the selection of sequencing technology should align with different stages of this information flow. This review includes various sequencing methods, such as WGS, RRBS, and RNA-seq. Each method has its technical advantages and limitations: WGS offers comprehensive genomic information but at a higher cost; RRBS is more cost-effective but provides a narrower scope of information; RNA-seq reveals changes in gene expression, which is crucial for understanding TED's molecular mechanisms. Mass spectrometry is a vital technique for high-throughput

sequencing in proteomics. For instance, MALDI-TOF serves as a cost-effective, efficient, and precise analytical tool widely used for microbial species identification and protein profiling in both clinical and research settings. MALDI-TOF can rapidly and accurately identify bacteria or proteins by analyzing their unique molecular fingerprints, thus providing valuable insights into their biological functions and characteristics [55-57]. Within the framework of the central dogma, combining transcriptomics and proteomics methods offers a comprehensive view from gene expression to changes at the protein and metabolite levels. This multi-omics approach elucidates the relationship between gene expression regulation and protein function, leading to a more thorough understanding of the molecular mechanisms underlying TED.

In the realm of TED research, the selection of an appropriate technological approach to address specific scientific inquiries is a critical step. Therefore, the choice of technology should be guided by specific research questions and available biological materials. For instance, while WGS provides comprehensive genomic information, not all studies require such exhaustive analysis [30]; RRBS or targeted gene panels might suffice. Similarly, for researchers interested in immediate molecular responses, RNA-seq may be more appropriate than microarrays. Proteomics can elucidate alterations in protein expression associated with inflammation, adipose tissue differentiation, extracellular matrix remodeling, and metabolism in the orbital tissues of TED patients [21]. Metabolomics can identify metabolic changes related to the pathogenesis of TED, such as alterations in cholesterol metabolism, which may be linked to the inflammatory and immune responses of the disease [24]. Although microbiomics may not be directly involved in the traditional central dogma, it provides a new perspective for understanding the systemic aspects of TED.

Multi-omics sequencing has identified a range of potential biomarkers for TED, offering promising applications for disease identification, course prediction, and treatment response evaluation. For example, Ji et al. explored the diagnostic value of a multi-panel approach combining various biomarkers to differentiate between TED and GD patients [58]. Their study demonstrated an area under the curve range of 0.845 to 0.935, suggesting that such multi-panel approaches could effectively identify individuals with TED. However, the clinical applicability of these findings depends on further validation in larger, more diverse cohorts to ensure their generalizability and reliability. Similarly, Zhang et al. used proteomic and miRNA analyses to identify 20 DEPs, including Zonulin,  $\alpha$ -2 macroglobulin,  $\beta$ -2 glycoprotein 1, and fibronectin, which may play roles in diagnosis and prognosis [59]. While clinical practice currently relies on markers such as TBII, TSHR-Ab, TSI and TSAb, additional potential biomarkers that could serve as adjuncts for TED confirmation and prognosis assessment remain to be identified and validated. Despite the progress made in biomarker identification through multi-omics sequencing, realizing their full clinical potential involves overcoming several challenges. Future research should focus on establishing well-defined conducting longitudinal studies, cohorts, and addressing technical, logistical, and economic barriers to the integration of these biomarkers into routine clinical practice.

Recent advancements in multi-omics technologies are enhancing the discovery and potential development of informative biomarkers for clinical practice [60]. These technologies offer valuable insights into the complex pathophysiology of TED and facilitate the identification of new diagnostic and therapeutic biomarkers [17, 21, 61]. Despite the promising prospects, several challenges remain, particularly regarding the translation of research discoveries into clinical applications. A significant issue is the inadequacy of study designs, which often results in low statistical significance and limits the impact of many research efforts. Future research in TED should continue to leverage multi-omics technologies, with a focus on improving sample collection, data standardization, and analytical methods [62]. Additionally, it is crucial to validate the clinical relevance of these biomarkers through larger sample sizes and multicenter studies to ensure their robustness and applicability in clinical settings.

It is important to recognize the limitations and challenges associated with the current body of research. Firstly, there is a need to increase sample sizes to enhance the robustness of findings. Larger sample sizes will improve the statistical power and generalizability of results. Secondly, standardization of biomarker detection methodologies across studies is crucial. Such standardization ensures the reliability and reproducibility of results, facilitating more accurate comparisons and interpretations. Thirdly, the validation of identified biomarkers in independent cohorts remains inadequately addressed. Many studies, according to the QUADOMICS criteria, have used patient populations that are not fully representative, and there has been insufficient validatory estimation and assessment, which may lead to overfitting. Fourthly, the lack of quantitative analysis prevents the performance of a meta-analysis, as the available data is not sufficiently homogeneous to draw meaningful conclusions or make reliable

statistical inferences. Future research should focus on overcoming these limitations by developing more representative patient cohorts, implementing standardized biomarker detection protocols, and rigorously validating biomarkers in independent studies. With continued research efforts, the integration of novel biomarkers holds the potential to significantly enhance disease management.

# Conclusion

This systematic review has explored the complex landscape of TED through the application of multi-omics sequencing, revealing a wide range of potential biomarkers that could significantly enhance diagnosis, monitoring, and treatment of this multifaceted condition. Despite variations in methodologies and the early stage of biomarker discovery, the evidence collectively highlights the transformative potential of genomics, transcriptomics, metabolomics, proteomics, lipidomics, and microbiomics unraveling the in intricate pathophysiology of TED. Our comprehensive analysis has demonstrated the value of diverse biological samples-such as orbital tissues, peripheral blood, feces, tears, and ocular microbiota-in providing insights into the molecular mechanisms of TED. These biomarkers reflect the systemic nature of the disease and underscore the potential for developing minimally invasive diagnostic approaches. The identification of upregulated genes and proteins associated with inflammation, adipogenesis, and immune dysregulation, along with unique metabolic and microbial signatures, presents promising avenues for future research of TED.

# **Supplementary Material**

Supplementary tables. https://www.ijbs.com/v20p6038s1.pdf

# Acknowledgements

## Funding

This work was supported by the National Natural Science Foundation of China (grant numbers 82388101, 81930024, and 82201235), the Science and Technology Commission of Shanghai (grant number 20DZ2270800), and the Shanghai Key Clinical Specialty, Shanghai Eye Disease Research Center (grant number 2022ZZ01003).

## **Consent for Publication**

All authors approved the final manuscript and the submission to this journal.

#### **Author Contributions**

HYZ: conception and design, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript for important intellectual content; YYZ: conception and design, acquisition and interpretation of data, drafting of the manuscript; BGY: analysis of data, drafting of the manuscript; YYD, YW, and SJF: analysis and interpretation of data, drafting of the manuscript; HFZ, XQF and XFS: critical revision of the manuscript for important intellectual content, administrative support, obtaining funding, supervision.

### **Competing Interests**

The authors have declared that no competing interest exists.

#### References

- 1. Bahn RS. Graves' ophthalmopathy. N Engl J Med. 2010; 362(8):726-38.
- Bartley GB. The epidemiologic characteristics and clinical course of ophthalmopathy associated with autoimmune thyroid disease in Olmsted County, Minnesota. Trans Am Ophthalmol Soc. 1994; 92:477– 588.
- Oculoplastic and Orbital Disease Group of Chinese Ophthalmological Society of Chinese Medical Association, Thyroid Group of Chinese Society of Endocrinology of Chinese Medical Association. [Chinese guideline on the diagnosis and treatment of thyroid-associated ophthalmopathy (2022)]. Zhonghua Yan Ke Za Zhi. 2022; 58(9):646–68.
- Bartalena L, Piantanida E, Gallo D, Lai A, Tanda ML. Epidemiology, Natural History, Risk Factors, and Prevention of Graves' Orbitopathy. Front Endocrinol. 2020; 11:615993.
- Hai YP, Saeed MEM, Ponto KA, Elflein HM, Lee ACH, Fang S, et al. A Multicenter, Single-Blind, Case-Control, Immunohistochemical Study of Orbital Tissue in Thyroid Eye Disease. Thyroid. 2022; 32(12):1547–58.
- Ross DS, Burch HB, Cooper DS, Greenlee MC, Laurberg P, Maia AL, et al. 2016 American Thyroid Association Guidelines for Diagnosis and Management of Hyperthyroidism and Other Causes of Thyrotoxicosis. Thyroid. 2016; 26(10):1343–421.
- Barrio-Barrio J, Sabater AL, Bonet-Farriol E, Velázquez-Villoria Á, Galofré JC. Graves' Ophthalmopathy: VISA versus EUGOGO Classification, Assessment, and Management. J Ophthalmol. 2015; 2015:249125.
- Fang S, Huang Y, Wang N, Zhang S, Zhong S, Li Y, et al. Insights into Local Orbital Immunity: Evidence for the Involvement of the Th17 Cell Pathway in Thyroid-Associated Ophthalmopathy. J Clin Endocrinol Metab. 2019; 104(5):1697–711.
- Huang Y, Wu Y, Zhang S, Lu Y, Wang Y, Liu X, et al. Immunophenotype of Lacrimal Glands in Graves' Orbitopathy: Implications for the Pathogenesis of Th1 and Th17 Immunity. Thyroid. 2022; 32(8):949–61.
- Verma R, Chen AJ, Choi D, Wilson DJ, Grossniklaus HE, Dailey RA, et al. Inflammation and Fibrosis in Orbital Inflammatory Disease: A Histopathologic Analysis. Ophthal Plast Reconstr Surg. 2023; 39(6):588– 93.
- Fang S, Huang Y, Wang S, Zhang Y, Luo X, Liu L, et al. IL-17A Exacerbates Fibrosis by Promoting the Proinflammatory and Profibrotic Function of Orbital Fibroblasts in TAO. J Clin Endocrinol Metab. 2016; 101(8):2955-65.
- Fang S, Huang Y, Zhong S, Zhang Y, Liu X, Wang Y, et al. IL-17A Promotes RANTES Expression, But Not IL-16, in Orbital Fibroblasts Via CD40-CD40L Combination in Thyroid-Associated Ophthalmopathy. Invest Ophth Vis Sci. 2016; 57(14):6123–33.
- Zhang H, Fang L, Cheng Y, Peng Y, Wang H, Jiang M, et al. Peripheral CD3\*CD4\* T cells as indicators of disease activity in thyroid eye disease: age-dependent significance. Graefes Arch Clin Exp Ophthalmol. 2024; 262(9):2985-2997.
- Akhoundova D, Rubin MA. Clinical application of advanced multi-omics tumor profiling: Shaping precision oncology of the future. Cancer Cell. 2022; 40(9):920–38.

- Zhang Y, Lee TY. Revealing the Immune Heterogeneity between Systemic Lupus Erythematosus and Rheumatoid Arthritis Based on Multi-Omics Data Analysis. Int J Mol Sci. 2022; 23(9):5166.
- Crick F. Central dogma of molecular biology. Nature. 1970; 227(5258):561–3.
- Wu L, Zhang S, Li X, Yao J, Ling L, Huang X, et al. Integrative transcriptomics and proteomic analysis of extraocular muscles from patients with thyroid-associated ophthalmopathy. Exp Eye Res. 2020; 193:107962.
- Chen MH, Liao SL, Chen MH, Tsou PL, Shih MJ, Chang TC, et al. Lysosome-Related Genes Are Regulated in the Orbital Fat of Patients with Graves' Ophthalmopathy. Invest Ophthalmol Vis Sci. 2008; 49(11):4760.
- Ezra DG, Krell J, Rose GE, Bailly M, Stebbing J, Castellano L. Transcriptome-level microarray expression profiling implicates IGF-1 and Wnt signalling dysregulation in the pathogenesis of thyroid-associated orbitopathy. J Clin Pathol. 2012; 65(7):608–13.
- Planck T, Shahida B, Parikh H, Ström K, Åsman P, Brorson H, et al. Smoking Induces Overexpression of Immediate Early Genes in Active Graves' Ophthalmopathy. Thyroid. 2014; 24(10):1524–32.
- Virakul S, Somparn P, Pisitkun T, van der Spek PJ, Dalm VASH, Paridaens D, et al. Integrative Analysis of Proteomics and DNA Methylation in Orbital Fibroblasts from Graves' Ophthalmopathy. Front Endocrinol. 2021; 11:619989.
- Du R, Wang F, Yang C, Hu J, Liu J, Jian Q, et al. Metabolic features of orbital adipose tissue in patients with thyroid eye disease. Front Endocrinol. 2023; 14:1151757.
- Cheng KC, Huang HH, Hung CT, Chen CC, Wu WC, Suen JL, et al. Proteomic analysis of the differences in orbital protein expression in thyroid orbitopathy. Graefes Arch Clin Exp Ophthalmol. 2013; 251(12):2777–87.
- Huang J, Chen M, Liang Y, Hu Y, Xia W, Zhang Y, et al. Integrative metabolic analysis of orbital adipose/connective tissue in patients with thyroid-associated ophthalmopathy. Front Endocrinol. 2022; 13:1001349.
- Virili C, Centanni M. Does microbiota composition affect thyroid homeostasis? Endocrine. 2015; 49(3):583–7.
- Virili C, Fallahi P, Antonelli A, Benvenga S, Centanni M. Gut microbiota and Hashimoto's thyroiditis. Rev Endocr Metab Disord. 2018; 19(4):293– 300.
- Biscarini F, Masetti G, Muller I, Verhasselt HL, Covelli D, Colucci G, et al. Gut Microbiome Associated with Graves' Disease and Graves' Orbitopathy: The INDIGO Multicenter European Study. J Clin Endocrinol Metab. 2023; 108(8):2065–77.
- Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. BMJ. 2021; 372:n160.
- Lumbreras B, Porta M, Márquez S, Pollán M, Parker LA, Hernández-Aguado I. QUADOMICS: an adaptation of the Quality Assessment of Diagnostic Accuracy Assessment (QUADAS) for the evaluation of the methodological quality of studies on the diagnostic accuracy of '-omics'-based technologies. Clin Biochem. 2008 Nov;41(16– 17):1316–25.
- Rotondo Dottore G, Bucci I, Lanzolla G, Dallan I, Sframeli A, Torregrossa L, et al. Genetic Profiling of Orbital Fibroblasts from Patients with Graves' Orbitopathy. J Clin Endocrinol Metab. 2021; 106(5):e2176-90.
- Liang Y, Ding S, Wang X, Hu C, Zhang Y, Hu Y, et al. Adipose/Connective Tissue from Thyroid-Associated Ophthalmopathy Uncovers Interdependence Between Methylation and Disease Pathogenesis: A Genome-Wide Methylation Analysis. Front Cell Dev Biol. 2021; 9:716871.
- Xin Z, Hua L, Shi TT, Tuo X, Yang FY, Li Y, et al. A genome-wide DNA methylation analysis in peripheral blood from patients identifies risk loci associated with Graves' orbitopathy. J Endocrinol Invest. 2018; 41(6):719–27.
- 33. Xin Z, Hua L, Yang YL, Shi TT, Liu W, Tuo X, et al. A Pathway Analysis Based on Genome-Wide DNA Methylation of Chinese Patients with Graves' Orbitopathy. Biomed Res Int. 2019; 2019:9565794.
- 34. Tao W, Ayala-Haedo JA, Field MG, Pelaez D, Wester ST. RNA-Sequencing Gene Expression Profiling of Orbital Adipose-Derived Stem Cell Population Implicate HOX Genes and WNT Signaling Dysregulation in the Pathogenesis of Thyroid-Associated Orbitopathy. Invest Ophthalmol Vis Sci. 2017; 58(14):6146-58.
- Bai XJ, Chu XJ, Yang T, Tong BD, Qi X, Peng YY, et al. Changes of mRNA expression underlying orbital adipogenesis in thyroid-associated orbitopathy. Arch Med Sci. 2022; 18(6):1708–15.
- Kim DW, Taneja K, Hoang T, Santiago CP, McCulley TJ, Merbs SL, et al. Transcriptomic Profiling of Control and Thyroid-Associated Orbitopathy (TAO) Orbital Fat and TAO Orbital Fibroblasts Undergoing Adipogenesis. Invest Ophthalmol Vis Sci. 2021; 62(9):24.

- Dorsky RI, Moon RT, Raible DW. Control of neural crest cell fate by the Wnt signalling pathway. Nature. 1998; 396(6709):370–3.
- Ross SE, Hemati N, Longo KA, Bennett CN, Lucas PC, Erickson RL, et al. Inhibition of adipogenesis by Wnt signaling. Science. 2000; 289(5481):950–3.
- 39. Wang D, Ling J, Tan R, Wang H, Qu Y, Li X, et al. CD169<sup>+</sup> classical monocyte as an important participant in Graves' ophthalmopathy through CXCL12-CXCR4 axis. iScience. 2024; 27(3):109213.
- Wang Y, Chen Z, Wang T, Guo H, Liu Y, Dang N, et al. A novel CD4<sup>+</sup> CTL subtype characterized by chemotaxis and inflammation is involved in the pathogenesis of Graves' orbitopathy. Cell Mol Immunol. 2021; 18(3):735-45.
- Chen Z, Zhang M, Liu Y, Chen Z, Wang L, Wang W, et al. VEGF-A enhances the cytotoxic function of CD4<sup>+</sup> cytotoxic T cells via the VEGF-receptor 1/VEGF-receptor 2/AKT/mTOR pathway. J Transl Med. 2023; 21:74.
- Kang J, Li Y, Zhao Z, Zhang H. Differentiation between thyroidassociated orbitopathy and Graves' disease by iTRAQ-based quantitative proteomic analysis. FEBS Open Bio. 2021; 11(7):1930–40.
- Ueland HO, Ueland GÅ, Løvås K, Breivk LE, Thrane AS, Meling Stokland AE, et al. Novel inflammatory biomarkers in thyroid eye disease. Eur J Endocrinol. 2022; 187(2):293–300.
- 44. Matheis N, Okrojek R, Grus FH, Kahaly GJ. Proteomics of Tear Fluid in Thyroid-Associated Orbitopathy. Thyroid. 2012; 22(10):1039–45.
- 45. Matheis N, Grus FH, Breitenfeld M, Knych I, Funke S, Pitz S, et al. Proteomics Differentiate Between Thyroid-Associated Orbitopathy and Dry Eye Syndrome. Invest Ophth Vis Sci. 2015; 56(4):2649.
- Kishazi E, Dor M, Eperon S, Oberic A, Hamedani M, Turck N. Thyroid-associated orbitopathy and tears: A proteomics study. J Proteomics. 2018; 170:110–6.
- Aass C, Norheim I, Eriksen EF, Børnick EC, Thorsby PM, Pepaj M. Comparative proteomic analysis of tear fluid in Graves' disease with and without orbitopathy. Clin Endocrinol. 2016; 85(5):805–12.
- Ji X, Dong K, Pu J, Yang J, Zhang Z, Ning X, et al. Comparison of the ocular surface microbiota between thyroid-associated ophthalmopathy patients and healthy subjects. Front Cell Infect Microbiol. 2022; 12:914749.
- Shi TT, Xin Z, Hua L, Zhao RX, Yang YL, Wang H, et al. Alterations in the intestinal microbiota of patients with severe and active Graves' orbitopathy: a cross-sectional study. J Endocrinol Invest. 2019; 42(8):967– 78.
- Shi TT, Xin Z, Hua L, Wang H, Zhao RX, Yang YL, et al. Comparative assessment of gut microbial composition and function in patients with Graves' disease and Graves' orbitopathy. J Endocrinol Invest. 2021; 44(2):297–310.
- Zhang Q, Tong B, Xie Z, Li Y, Li Y, Wang L, et al. Changes in the gut microbiota of patients with Graves' orbitopathy according to severity grade. Clin Exp Ophthalmol. 2023; 51(8):808–21.
- Shi TT, Hua L, Wang H, Xin Z. The Potential Link between Gut Microbiota and Serum TRAb in Chinese Patients with Severe and Active Graves' Orbitopathy. Int J Endocrinol. 2019; 2019:1–12.
- 53. Fenneman AC, van der Spek AH, Hartstra A, Havik S, Salonen A, de Vos WM, et al. Intestinal permeability is associated with aggravated inflammation and myofibroblast accumulation in Graves' orbitopathy: the MicroGO study. Front Endocrinol. 2023; 14:1173481.
- Billiet B, Chao de la Barca JM, Ferré M, Muller J, Vautier A, Assad S, et al. A Tear Metabolomic Profile Showing Increased Ornithine Decarboxylase Activity and Spermine Synthesis in Thyroid-Associated Orbitopathy. J Clin Med. 2022; 11(2):404.
- Croxatto A, Prod'hom G, Greub G. Applications of MALDI-TOF mass spectrometry in clinical diagnostic microbiology. FEMS Microbiol Rev. 2012; 36(2):380–407.
- Hajduk J, Matysiak J, Kokot ZJ. Challenges in biomarker discovery with MALDI-TOF MS. Clin Chim Acta. 2016; 458:84–98.
- Patel R. MALDI-TOF MS for the diagnosis of infectious diseases. Clin Chem. 2015; 61(1):100–11.
- Ji DY, Park SH, Park SJ, Kim KH, Ku CR, Shin DY, et al. Comparative assessment of Graves' disease and main extrathyroidal manifestation, Graves' ophthalmopathy, by non-targeted metabolite profiling of blood and orbital tissue. Sci Rep. 2018; 8:9262.
- Zhang L, Masetti G, Colucci G, Salvi M, Covelli D, Eckstein A, et al. Combining micro-RNA and protein sequencing to detect robust biomarkers for Graves' disease and orbitopathy. Sci Rep. 2018; 8:8386.
- Li C, Sun YD, Yu GY, Cui JR, Lou Z, Zhang H, et al. Integrated Omics of Metastatic Colorectal Cancer. Cancer Cell. 2020; 38(5):734-747.e9.
- 61. Shi TT, Liu HY, Zhu XR, Xin Z, Hua L, Xie RR, et al. Integrated Proteomics and Metabolomics Analyses of Serum in Chinese Patients with Severe and Active Graves' Orbitopathy: A Cross-sectional Study. Endocr Metab Immune. 2023; 23(9):1151–61.

- Xiao Y, Bi M, Guo H, Li M. Multi-omics approaches for biomarker discovery in early ovarian cancer diagnosis. EBioMedicine. 2022; 79:104001.
- Zhu L, Li S, He S, Tong Q, Wang L, Li X, et al. The critical role of m6A methylation in the pathogenesis of Graves' ophthalmopathy. Eye Vis (Lond). 2020; 7:55.
- 64. Sun A, Ye H, Xu Z, Chen J, Xiao W, Zhang T, et al. Serelaxin Alleviates Fibrosis in Thyroid-Associated Ophthalmopathy via the Notch Pathway. Int J Mol Sci. 2023 May 6;24(9):8356.
- 65. Ye H, Sun A, Xiao W, Zhang T, Xu Z, Shi L, et al. Differential Circular RNA Expression Profiling of Orbital Connective Tissue from Patients with Type I and Type II Thyroid-Associated Ophthalmopathy. Invest Ophth Vis Sci. 2022; 63(12):27.
- 66. Wu L, Liang Y, Song N, Wang X, Jiang C, Chen X, et al. Differential expression and alternative splicing of transcripts in orbital adipose/connective tissue of thyroid-associated ophthalmopathy. Exp Biol Med. 2021; 246(18):1990–2006.
- Khong JJ, Wang LY, Smyth GK, McNab AA, Hardy TG, Selva D, et al. Differential Gene Expression Profiling of Orbital Adipose Tissue in Thyroid Orbitopathy. Invest Ophth Vis Sci. 2015; 56(11):6438–47.
- Wu L, Zhou R, Diao J, Chen X, Huang J, Xu K, et al. Differentially expressed circular RNAs in orbital adipose/connective tissue from patients with thyroid-associated ophthalmopathy. Exp Eye Res. 2020; 196:108036.
- Wang Y, Liu Y, Cai J, Zong T, Zhang Z, Xie T, et al. Differentially expressed genes in orbital adipose/connective tissue of thyroid-associated orbitopathy. PeerJ. 2023; 11:e16569.
- Wu L, Li L, Liang Y, Chen X, Mou P, Liu G, et al. Identification of differentially expressed long non-coding RNAs and mRNAs in orbital adipose/connective tissue of thyroid-associated ophthalmopathy. Genomics. 2021; 113(1):440–9.
- Mou P, Chen Z, Jiang L, Cheng J, Wei R. PTX3: A Potential Biomarker in Thyroid Associated Ophthalmopathy. Biomed Res Int. 2018; 2018;5961974.
- Li Z, Wang M, Tan J, Zhu L, Zeng P, Chen X, et al. Single-cell RNA sequencing depicts the local cell landscape in thyroid-associated ophthalmopathy. Cell Rep Med. 2022; 3(8):100699.
- Lee BW, Kumar VB, Biswas P, Ko AC, Alameddine RM, Granet DB, et al. Transcriptome Analysis of Orbital Adipose Tissue in Active Thyroid Eye Disease Using Next Generation RNA Sequencing Technology. Open Ophthalmol J. 2018; 12:41–52.
- Planck T, Parikh H, Brorson H, Mårtensson T, Åsman P, Groop L, et al. Gene Expression in Graves' Ophthalmopathy and Arm Lymphedema: Similarities and Differences. Thyroid. 2011; 21(6):663–74.
- Lantz M, Vondrichova T, Parikh H, Frenander C, Ridderstråle M, Åsman P, et al. Overexpression of Immediate Early Genes in Active Graves' Ophthalmopathy. J Clin Endocrinol Metab. 2005; 90(8):4784–91.
- Rosenbaum JT, Choi D, Wong A, Wilson DJ, Grossniklaus HE, Harrington CA, et al. The Role of the Immune Response in the Pathogenesis of Thyroid Eye Disease: A Reassessment. PLoS One. 2015; 10(9):e0137654.
- Ma R, Gan L, Guo J, Peng Z, Wu J, Harrison AR, et al. Insights into Ferroptosis: Targeting Glycolysis to Treat Graves' Orbitopathy. J Clin Endocrinol Metab. 2022; 107(7):1994–2003.
- Jang SY, Chae MK, Lee JH, Lee EJ, Yoon JS. Role of miR-146a in the Regulation of Inflammation in an In Vitro Model of Graves' Orbitopathy. Invest Ophth Vis Sci. 2016; 57(10):4027–34.
- Kumar S, Leontovich A, Coenen MJ, Bahn RS. Gene expression profiling of orbital adipose tissue from patients with Graves' ophthalmopathy: a potential role for secreted frizzled-related protein-1 in orbital adipogenesis. J Clin Endocrinol Metab. 2005; 90(8):4730–5.
- Ko J, Kim YJ, Choi SH, Lee CS, Yoon JS. Yes-Associated Protein Mediates the Transition from Inflammation to Fibrosis in Graves' Orbitopathy. Thyroid. 2023; 33(12):1465–75.
- Yue Z, Tong F, Zeng C, Wei R. Identification of tRNA-Related Fragments and Their Potential Regulatory Effects in Thyroid-Associated Ophthalmopathy. Front Genet. 2022; 13:878405.
- Yue Z, Mou P, Chen S, Tong F, Wei R. A Novel Competing Endogenous RNA Network Associated with the Pathogenesis of Graves' Ophthalmopathy. Front Genet. 2021; 12:795546.
- Kim N, Choung H, Kim YJ, Woo SE, Yang MK, Khwarg SI, et al. Serum microRNA as a potential biomarker for the activity of thyroid eye disease. Sci Rep. 2023; 13:234.
- Liu R, Ye Z, Liu Q, Xuan M, Li R, Zhang L, et al. MicroRNA-376b is involved in the pathogenesis of thyroid-associated ophthalmopathy by regulating HAS2. Endocrine. 2023; 82(1):87–95.
- 85. Wescombe L, Lahooti H, Gopinath B, Wall JR. The cardiac calsequestrin gene (CASQ2) is up-regulated in the thyroid in patients with Graves'

ophthalmopathy – support for a role of autoimmunity against calsequestrin as the triggering event. Clin Endocrinol. 2010; 73(4):522–8.

- Matheis N, Lantz M, Grus FH, Ponto KA, Wolters D, Brorson H, et al. Proteomics of Orbital Tissue in Thyroid-Associated Orbitopathy. J Clin Endocrinol Metab. 2015; 100(12):E1523–30.
- Shi TT, Zhao RX, Xin Z, Hou ZJ, Wang H, Xie RR, et al. Tear-derived exosomal biomarkers of Graves' ophthalmopathy. Front Immunol. 2022; 13:1088606.
- Zhou X, Wei R, Wang R. Characterization of human tear proteome reveals differentially abundance proteins in thyroid-associated ophthalmopathy. PeerJ. 2022; 10:e13701.
- Jiang L, Wei R, Diao J, Ding H, Wang W, Ao R. Proteomics of Tear in Inactive Thyroid-Associated Ophthalmopathy. Acta Endocrinol (Buchar). 2021; 17(3):291–303.
- Chng CL, Seah LL, Yang M, Shen SY, Koh SK, Gao Y, et al. Tear Proteins Calcium binding protein A4 (S100A4) and Prolactin Induced Protein (PIP) are Potential Biomarkers for Thyroid Eye Disease. Sci Rep. 2018; 8:16936.
- 91. Okrojek R, Grus FH, Matheis N, Kahaly GJ. Proteomics in Autoimmune Thyroid Eye Disease. Horm Metab Res. 2009; 41(06):465–70.
- Byeon SK, Park SH, Lee JC, Hwang S, Ku CR, Shin DY, et al. Lipidomic differentiation of Graves' ophthalmopathy in plasma and urine from Graves' disease patients. Anal Bioanal Chem. 2018; 410(27):7121–33.
- Ueland HO, Ulvik A, Løvås K, Wolff ASB, Breivik LE, Stokland AEM, et al. Systemic Activation of the Kynurenine Pathway in Graves' Disease with and without Ophthalmopathy. J Clin Endocrinol Metab. 2023; 108(6):1290–7.
- 94. Li Y, Wang F, Hou Z, Nie Z, Ma L, Hui S, et al. Microbiome in orbital fat under thyroid associated ophthalmopathy. Medicine in Microecology. 2022; 13:100058.